

| Report Documentation Page | | | | Form Approved OMB No. 0704-0188 | |
|--|------------------------------------|-------------------------------------|--|--|---------------------------------|
| Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. | | | | | |
| 1. REPORT DATE 29 MAR 2011 | | 2. REPORT TYPE research | | 3. DATES COVERED 01-01-2010 to 01-01-2011 | |
| 4. TITLE AND SUBTITLE Specific recognition and detection of MRSA based on molecular probes comprised of lytic phage and antibody | | | | 5a. CONTRACT NUMBER | |
| | | | | 5b. GRANT NUMBER | |
| | | | | 5c. PROGRAM ELEMENT NUMBER | |
| 6. AUTHOR(S) | | | | 5d. PROJECT NUMBER | |
| | | | | 5e. TASK NUMBER | |
| | | | | 5f. WORK UNIT NUMBER | |
| 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Auburn University, College of Veterinary Medicine, Auburn, AL, 36849 | | | | 8. PERFORMING ORGANIZATION REPORT NUMBER | |
| 9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) 81st Medical Group, Clinical Research Laboratory, 301 Fisher Street, Keesler AFB, MS, 39534 | | | | 10. SPONSOR/MONITOR'S ACRONYM(S) 81 MDG, CRL | |
| | | | | 11. SPONSOR/MONITOR'S REPORT NUMBER(S) | |
| 12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited | | | | | |
| 13. SUPPLEMENTARY NOTES | | | | | |
| 14. ABSTRACT A Q-sense E4 QCM system was employed to study bacteria-phage interactions. | | | | | |
| 15. SUBJECT TERMS | | | | | |
| 16. SECURITY CLASSIFICATION OF: | | | 17. LIMITATION OF ABSTRACT 1 | 18. NUMBER OF PAGES 1 | 19a. NAME OF RESPONSIBLE PERSON |
| a. REPORT unclassified | b. ABSTRACT unclassified | c. THIS PAGE unclassified | | | |

Specific recognition and detection of MRSA based on molecular probes comprised of lytic phage and antibody

Rajesh Guntupalli^{*1}, Eric Olsen,² Ludmila Globa¹, Jacob Bosarge¹, Timothy Moore¹, Iryna Sorokulova¹, and Vitaly Vodyanoy¹

¹ College of Veterinary Medicine, Auburn University, Auburn, AL 36849. ² Clinical Research Laboratory, 81st Medical Group, Keesler AFB MS USA 39534

Methicillin resistant strains of *Staphylococcus aureus* (MRSA) are implicated in serious infections and nosocomial outbreaks. MRSA show resistance to a wide range of antibiotics thus limiting the treatment options. Therefore, rapid detection of MRSA is of key importance in prevention and diagnosis of infections caused by antibiotic resistant *S. aureus*. Currently existing methods of MRSA detection have some limitations and lack sensitivity or specificity. We examined a new approach in specific recognition and detection of MRSA, including identification of bacteria together with conformation of MRSA in real time. For this purpose we use a newly isolated *S. aureus* bacteriophage with a wide spectrum of hosts (including MRSA strains) together with monoclonal antibody against a penicillin-binding protein (PBP 2a). PBP 2a is a cell wall protein and it is responsible for antibiotic resistivity of MRSA. We showed that simultaneous recognition of *Staphylococcus* bacteria and PBP 2a protein increases specificity and reliability of MRSA detection.

A Q-sense E4 QCM-D system (Sweden) was employed to study bacteria-phage interactions. Lytic phages were constructed into hollow spherical particles upon exposure to a chloroform-water interface. These particles were converted into monolayers and deposited onto QCM-D crystals using Langmuir-Blodgett technique [1]. Binding of MRSA has been manifested by frequency and dissipation changes (Fig.1). Similar results were obtained with antibiotic sensitive *S. aureus*.

An agglutination test was carried out using a latex reagent sensitized with antibody against PBP 2a (Denka Seiken Co., Ltd, Tokyo, Japan). The bacterial cells were mixed with the latex reagent on a test card. A visible agglutination occurred in about three minutes for MRSA. For the antibiotic sensitive *S. aureus* strains no agglutination occurred, even after one hour.

The interaction between PBP 2a protein and antibodies is not specific for *Staphylococcus aureus* because other bacteria have antibiotic binding proteins with large sequence similarity to PBP 2a. Therefore in our work we used *S. aureus* bacteriophage with wide spectrum of hosts together with antibodies against PBP 2a protein. In order to build a biosensor to specifically detect and identify MRSA we will employ a device with two parallel channels. One channel will have a *S. aureus* bacteriophage monolayer as a sensor probe, while the sensor of another channel will be covered with PBP 2a specific antibodies. Consequently, one channel will identify *S. aureus* bacteria, while another one will be sensitive to the antibiotic-binding protein. When signals coming from two channels are positive it would indicate the specific detection of MRSA.

Acknowledgements Supported by grants from USAF CRADA 07-277-60MDG-01 and Auburn University AUDFS. The views expressed in this article are those of the authors, and do not reflect the official policy or position of the United States Air Force, Department of Defense, or the U.S. Government.

[1] Guntupalli, R, Sorokulova, I, Long, R, Olsen, E, Neely, W, Vodyanoy, V (2010) Phage Langmuir monolayers and Langmuir-Blodgett films. *Colloids and Surfaces B-Biointerfaces* 82: 182-189.




Figure 1. Time change in frequency and dissipation energy during MRSA binding